

LARGE-SCALE DENSITY–VELOCITY RELATIONS

M.J. CHODOROWSKI

Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland



I present recent progress in theoretical modelling of cosmological density–velocity relations in the weakly nonlinear regime. The relations are local, based on rigorous perturbation theory and include the effects of smoothing of the density and the velocity fields. For small smoothing scales, they can be improved by slight adjustments based on N-body results. The relations can be useful for density–velocity comparisons, yielding a fair estimate of Ω and offering a method for disentangling Ω and bias.

1 Introduction

In the gravitational instability paradigm for the formation of structure in the Universe, the peculiar motions of galaxies are tightly related to the large-scale mass distribution. The comparison between the density and the velocity fields can serve as a test of the gravitational instability hypothesis and as a method for estimating the cosmological parameter Ω (Dekel *et al.*⁷). In linear theory, the relation between the density and the velocity fields is

$$\delta(\mathbf{x}) = -f(\Omega)^{-1} \nabla \cdot \mathbf{v}(\mathbf{x}), \quad (1)$$

where $f(\Omega) \simeq \Omega^{0.6}$. The relation between the density contrast and the velocity divergence is thus linear and local.

This equation is applicable only when the density fluctuations are small compared to unity, $\sigma_\delta^2 \equiv \langle \delta^2 \rangle \ll 1$. However, sampling of galaxies in current redshift surveys and random errors in the peculiar velocity data enable reliable dynamical analysis at smoothing scales down to a few h^{-1} Mpc, where fluctuations already exceed the regime of applicability of linear theory. On the other hand, even at scales as small as a few h^{-1} Mpc, ‘typical’ (rms) fluctuations are still not in excess of unity.^a One could thus hope that to construct the relation between the fields in question, perturbation theory can be effectively used.

In perturbation theory, one approximates the evolved density contrast as a sum of terms δ_j , each corresponding to the j^{th} order in perturbation theory,

$$\delta = \delta_1 + \delta_2 + \delta_3 + \dots \quad (2)$$

Similarly, defining a variable θ proportional to the velocity divergence,

$$\theta = -f(\Omega)^{-1} \nabla \cdot \mathbf{v}, \quad (3)$$

we can expand it in a perturbative series,

$$\theta = \theta_1 + \theta_2 + \theta_3 + \dots \quad (4)$$

^aFor a Gaussian smoothing function, commonly used in the density–velocity comparisons.

The solutions for δ_j and θ_j were obtained more than a decade ago (Fry⁸, Goroff *et al.*⁹). In general, the j^{th} order solution is of the order of $(\delta_1)^j \sim \sigma_\delta^j$. In the weakly nonlinear regime, i.e., when $\sigma_\delta < 1$, which is our case, we may hope the perturbative series to converge rapidly.

Linear theory is nothing but perturbation theory truncated at the lowest, i.e. the first order. A natural way of extending the linear relation (1) is to include higher-order contributions to the density contrast and the velocity divergence. Unfortunately, these contributions are non-local, hence the nonlinear density vs velocity-divergence relation (DVDR) *at a given point* has a scatter. Therefore, as a local estimator of the density from the velocity divergence we adopt the conditional mean, i.e., the mean density *given* the velocity divergence.

The calculation of the conditional moments is a statistical problem. To compute mean δ given θ , $\langle \delta \rangle | \theta$, we need the joint probability distribution function (PDF) $p(\delta, \theta)$. Under an assumption of Gaussian initial conditions, adopted in this work, this function is initially a bivariate Gaussian (degenerated, since $\delta_1 = \theta_1$). Subsequent nonlinear gravitational evolution, however, drives the PDF away from its initial shape. Our problem here is to calculate it in the weakly nonlinear regime.

Mathematically, a PDF is given by the inverse Fourier transform of its characteristic function Φ , which in turn is related to the cumulant generating function $\mathcal{K}$ by the equation $\Phi = \exp[\mathcal{K}]$. The cumulants, κ_{mn} , from which $\mathcal{K}$ is constructed,

$$\mathcal{K}(it, is) = \sum_{(m,n) \neq (0,0)}^{\infty} \frac{\kappa_{mn}}{m!n!} (it)^m (is)^n, \quad (5)$$

are the *connected* (reduced) part of the joint moments of δ and θ , $\kappa_{mn} = \langle \delta^m \theta^n \rangle_{\text{conn}}$. We introduce $\mathcal{K}$ because in the weakly nonlinear regime, the cumulants obey a scaling hierarchy in σ_δ such that the series (5) is a power series in σ_δ (even in the case of the standard variables $\mu \equiv \delta/\sigma_\delta$ and $\nu \equiv \theta/\sigma_\theta$). Therefore, truncating (5) at order p we neglect contributions which are $\mathcal{O}(\sigma_\delta^{p+1})$. This yields the truncated PDF and in turn the weakly nonlinear DVDR up to a given order in perturbation theory.

2 Results

Up to third order (Chodorowski & Łokas³) we have

$$\langle \delta \rangle | \theta = a_1 \theta + a_2 (\theta^2 - \sigma_\theta^2) + a_3 \theta^3, \quad (6)$$

where σ_θ^2 is the variance of the velocity divergence field. The reverse relation (Chodorowski *et al.*⁵) is

$$\langle \theta \rangle | \delta = r_1 \delta + r_2 (\delta^2 - \sigma_\delta^2) + r_3 \delta^3 \quad (7)$$

(an extension of a second-order formula by Bernardeau¹). Because the DVDR has a scatter, expression (7) cannot in general be obtained by direct inversion of expression (6). In density–density comparisons, one reconstructs the mass density field from the observed velocity field and compares it to the observed galaxy density field. In velocity–velocity comparisons, one reconstructs (under some assumptions about bias) the velocity field from the observed galaxy density field and compares to the observed velocity field. Formula (6) is thus relevant for density–density comparisons, while formula (7) is relevant for velocity–velocity comparisons. Indeed, for irrotational flows the velocity field can readily be recovered from its divergence given some boundary conditions at large distances,

$$\mathbf{v}(\mathbf{x}) = \frac{f(\Omega)}{4\pi} \int d^3x' \theta(\mathbf{x}') \frac{\mathbf{x}' - \mathbf{x}}{|\mathbf{x}' - \mathbf{x}|^3}. \quad (8)$$




Figure 1: Left: joint probability distribution function for the density contrast (δ) and the velocity divergence (θ) from an N-body simulation by Chodorowski *et al.*,⁶ shown against $1 + \delta$ and $1.5 + \theta$. Right: the coefficients a_j (see eq. [6]), as functions of the smoothing scale, from N-body simulations by Chodorowski & Stompor.⁴ Simulation results are shown as thin curves; third order perturbation theory predictions are shown as thick lines. The simulations are for power-law power spectra with the spectral index n equal to -1.5 (left) and -1 (right). A top-hat smoothing function is used.

The coefficients a_j and r_j are the previrialization-, skewness-, and kurtosis-like combinations of the joint moments of δ and θ . Calculations of the reduced moments have already become a small industry in perturbation theory. For Gaussian smoothing, the related moments for δ and for θ (separately) were computed by Łokas *et al.*¹⁰ and Łokas *et al.*¹¹ The coefficients a_j and r_j , though constructed from the joint moments, possess similar mathematical structure and were computed in an analogous way.^{3,5}

Chodorowski *et al.*⁵ also calculated the scatter in the DVDR and found it to be relatively small.^b Nevertheless, one can try to further reduce the variance of density estimators by including off-diagonal components of the velocity deformation tensor. Indeed, including the shear of the velocity field reduces the scatter in the density–velocity relation.^{12,2}

3 Tests against N-body simulations

Perturbative expressions for density–velocity relations should be checked against N-body simulations, in order to assess their range of applicability. The results of the simulations by Chodorowski *et al.*⁶ and Chodorowski & Stompor⁴ (see Figure 1) can be summarized as follows:

- For large smoothing scales ($\sigma_\delta \ll 1$), the numerical coefficients a_j and r_j converge to the predicted values. (In this sense, the predicted DVDRs are ‘asymptotically unbiased’.)
- For smoothing scales as small as a few megaparsecs ($\sigma_\delta \lesssim 1$),
 - The numerical coefficients slightly, but systematically, deviate from predicted values. (Apparently, higher–than third–order contributions become significant.)
 - On the other hand, third-order polynomials, with fitted coefficients, provide excellent fits to N-body data. (Higher-order contributions modify the values of the coefficients for $j = 1, 2, 3$, but do not significantly induce non-zero values for $j > 3$.)

^bThe ratio of the scatter to the rms fluctuation of the density field, σ_δ , vanishes in the limit $\sigma_\delta \rightarrow 0$, as expected.

4 Implications for the value of Ω

For the sake of simplicity, here we will consider only second-order corrections to the linear relation (1). Neglecting third-order terms in equation (7) yields

$$\theta = \delta - a_2(\delta^2 - \sigma_\delta^2) \quad (9)$$

($r_2 = -a_2$, $0.2 < a_2 < 0.3$ ^{5,3}). Assuming linear bias between the galaxy density field, δ_g , and the mass density field, $\delta_g = b\delta$, we can express the relation in terms of the observable quantities $\nabla \cdot \mathbf{v}$ and δ_g ,

$$-\nabla \cdot \mathbf{v} = \beta\delta_g - a_2b^{-1}\beta\delta_g^2. \quad (10)$$

Here, $\beta(\Omega, b) \equiv f(\Omega)/b$ and we also neglect the offset in the relation. We see that the quadratic correction has the opposite sign to the linear term. It implies that the linear relation overestimates the predicted velocities. Hence, in order to match the observed ones, a lower value of β is needed. In effect, *applying linear theory biases the estimate of Ω low* (cf. Strauss & Willick¹³).

Note that the two coefficients of the binomial in δ_g on the RHS of equation (10) are different combinations of β and b , or $f(\Omega)$ and b . This offers a way to solve for Ω and b separately.³

5 Summary

The weakly nonlinear estimators of density from velocity, and vice versa, presented here,

- are asymptotically unbiased;
- for smaller smoothing scales, can be improved by slight adjustments from N-body data;
- should be used in density–density and velocity–velocity comparisons, in order not to bias the estimate of Ω low;
- offer a method for disentangling Ω and bias.

Acknowledgments

This research has been supported in part by the Polish State Committee for Scientific Research grants No. 2.P03D.008.13 and 2.P03D.004.13.

References

1. F. Bernardeau, ApJ **390**, L61 (1992).
2. M.J. Chodorowski, MNRAS **292**, 695 (1997).
3. M.J. Chodorowski and E.L. Lokas, MNRAS **287**, 591 (1997).
4. M.J. Chodorowski and R. Stompor, in preparation.
5. M.J. Chodorowski, E.L. Lokas, A. Pollo and A. Nusser, preprint astro-ph/9802050.
6. M.J. Chodorowski, E.L. Lokas, F. Bernardeau and A. Kudlicki, in preparation.
7. A. Dekel, E. Bertschinger, A. Yahil, M. Strauss, M. Davis and J. Huchra, ApJ **412**, 1 (1993).
8. J.N. Fry, ApJ **279**, 499 (1984).
9. M.H. Goroff, B. Grinstein, S.-J. Rey and M.B. Wise, ApJ **311**, 6 (1986).
10. E.L. Lokas, R. Juszkiewicz, D.H. Weinberg and F.R. Bouchet, MNRAS **274**, 730 (1995).
11. E.L. Lokas, R. Juszkiewicz, F.R. Bouchet and E. Hivon, ApJ **467**, 1 (1996).
12. P.J. Mancinelli and A. Yahil, ApJ **452**, 75 (1995).
13. M.A. Strauss and J.A. Willick, Phys. Rep. **261**, 271 (1995).